

Production of four leptons in electroweak $\gamma\gamma$ interactions

I. Sotsky

NC PHEP ,BSU, Minsk, Belarus

Abstract

Detailed numerical analysis of four charged leptons photoproduction in frame of standard theory of electroweak interaction is presented. Total and differential cross sections are obtained and investigated using the Monte-Carlo method of numerical integration. Different energies of initial photons (60-2000 GeV in c.m.s.) as well as the definite and averaged spin states of interacting particles are considered. A set of kinematical cuts of linear collider experiments is discussed.

1 Introduction

There are planned to realize high energy experiments on linear collider which will have an opportunity to operate in $\gamma\gamma$, γe and ee mode [1]. This capability will provide new advantages in investigation of new physics, new particles research and detailed study of non abelian nature of electroweak interaction. However it is necessary to take into account several important features to realize such experiments successfully.

First of all, since produced $W^\pm$ and Higgs particles ordinarily decay within detector they could be observed via it's decay products only, for example via several pairs of leptons in final state. Because of high accuracy and relatively clean environment provided by leptonic collider, the exact calculation of all backgrounds, namely $\gamma\gamma \rightarrow 2l$, $\gamma\gamma \rightarrow 2l + \textit{photons}$ and $\gamma\gamma \rightarrow 4l$ processes, are required to study of such productions.

Secondly, since initial high energy γ quanta are produced by Compton backscattering of laser ray on fast electrons, the exact information about polarization state and energy spectrum of obtained photon's beams is necessary to interpret the results of such experiments correctly. This data can be gathered from measuring of several calibration processes, for example: $\gamma\gamma \rightarrow 2l$, $\gamma\gamma \rightarrow 2l + \textit{photons}$ and $\gamma\gamma \rightarrow 4l$.

So precision investigation of processes listed behind has great importance.

The analysis of $\gamma\gamma \rightarrow 2l$ and $\gamma\gamma \rightarrow 2l + \textit{photons}$ processes one can find in refs. [2]-[4]. The detailed theoretical description of $\gamma\gamma \rightarrow 4l$ reaction, a lot of useful references to the papers devoted to same problem and expression for squared matrix element constructed by helicity amplitude method one can find in ref. [5]. At this paper we perform numerical analysis of two charged leptons pair photoproduction in frame of Standard Model (SM) of electroweak interaction. A set of interacting energies, polarization states and experimental kinematical cuts are considered and discussed.

2 Calculation

There are six topologically different Feynman diagrams describing process of $\gamma\gamma \rightarrow 4l$ at the lowest order of SM (see fig.1). Whole set of diagrams can be derived on base of these six ones using C- P- and crossing symmetries.




Fig.1. Feynman diagrams for process $\gamma\gamma \rightarrow 4l$.

The diagrams containing charged current exchange are excluded because only processes with four charged leptons in the final state are considered.

Corresponding cross section has the form:

$$\sigma = \frac{1}{4(k_1 k_2)} \int |M|^2 d\Gamma, \quad (1)$$

where

$$d\Gamma = \frac{d^3 p_1}{(2\pi)^3 2p_1^0} \frac{d^3 p_2}{(2\pi)^3 2p_2^0} \frac{d^3 p_3}{(2\pi)^3 2p_3^0} \frac{d^3 p_4}{(2\pi)^3 2p_4^0} (2\pi)^4 \delta(k_1 + k_2 - p_1 - p_2 - p_3 - p_4)$$

is phase space element, p_i - four momentum of final lepton and k_i - four momentum of initial photon and (k_1, k_2) - scalar product of two four vectors.

The construction of squared matrix element is realized both using helicity amplitude method [6]-[9] as well as precision covariant one (see, for example, refs [10],[11]). Helicity amplitude method is used to obtain cross sections in massless limit at each possible polarization state configuration of interacting particles. Due to compact form of final expression it is allowed to perform numerical integration with high enough accuracy for a short interval of computing time. The explicit matrix element expression constructed by using helicity amplitude method one can find in ref. [5].

The covariant method of precision calculation allows to write squared matrix element without any approximation. It is applied to estimate error of massless approximation and to investigate heavy leptons production processes. The last method was used for calculation of cross section in case of unpolarized particles interaction.

The investigation of differential and total cross sections is realized by using of the Monte-Carlo method of numerical integration.

3 Results and Conclusion

We have calculated the differential and total cross sections of $\gamma\gamma \rightarrow 4l$ process in frame of SM at set of initial particles energies, polarization states and kinematical cuts.

The values of total cross sections at different kinematical conditions are presented in the Table. The differential cross section dependence on cosine of polar angle (angle between final lepton and initial photon) is shown at figs. 2-9. The case of averaged polarization state of interacting particles and fixed one are considered.

Presented results demonstrate the expected dependence of cross section on cosine of polar angle: cross section is almost equal to zero at the center of kinematical region (the kinematical region where final particles have large value of $p_{\perp}$) and extremely increases if one closed to the borders (the kinematical region with small $p_{\perp}$) (see figs. 2-9). The total cross section also strongly increases with decreasing of interacting photons energy (see the Table and figs. 14,15), owing to its expression contains $\frac{1}{k_1 k_2}$ factor, where k_1 and k_2 are four momenta of initial photons.

The polarized assymetry of differential cross section (figs. 4,5) occurs because of the probability of lepton scattering into direction closed to momentum of initial photon is larger in case of different signs of photon polarization and scattering lepton helicity then in case of the same signs.

As one can see at figs. 6,7 the differential cross section dependencies are symmetric since the initial photons have the similar polarization state, but the cross section value is larger when the sign of lepton helicity is opposite to polarization signs of interacting photons. The total cross section of lepton photoproduction with fixed polarization states is larger in case of different polarizations of initial photos then in case of similar ones (see figs. 8,9).

At figs. 10-13 the differential cross section dependencies on a set of kinematics variables are presented. Peaks which one can see at fig.10 are so colled collinear peaks. This peaks appear when several particles propagate into very closed direction. Cross section incidence near borders of kinimatical region is caused by application of kinematical cuts applying. Peak at fig. 11 is well noticeable so called Z-peak, which is appeared if mass of any final pair is closed to mass of Z-boson. At fig. 12 one can see that cross section extremely increases when energies of two final electron positron pairs are closed to each other. Fig. 13 demonstrates that cross section has the largest value when any final particle has the minimal (or the maximal) admissible value of energy.

The comparison of results obtained by the helicity amplitude method and the precision covariant one indicates that electron and positron mass contributions are vanishingly small under investigated kinematical condition: energy of initial photons beams is $60 \sim 2000$ GeV, polar angle is larger then 7° , and angle between any two final particles is larger then 3° .

Besides that, as one can see in the Table, the distinguish between cross sections of processes electrons and muons photoproduction becomes less then statistical error of Monte-Carlo method when energy of interacting particles equals to 0.5 TeV in c.m.s. So if the energy of initial photons is equal to or higher then 0.5 TeV in c.m.s the muon mass contribution could be neglected in frame of precision of planned experiments. However, it is imposible to neglect τ lepton mass contribution anywhere at investigated kinematical field (see the Table).

The relative error of obtained results ($\sim 0.9\%$) is less then expected experimental error in future high energy experiments on linear collider [1].

We have obtained the differential and total cross sections in frame of precision covariant method for unpolarized interacting particles only, so it is impossible to use them for investigation of polarized leptons production processes. However, it was discovered that one can perform spin effects analyzing in $\gamma\gamma \rightarrow \mu^+\mu^-\mu^+\mu^-$ and $\gamma\gamma \rightarrow e^+e^-\mu^+\mu^-$ reactions with enough precision

applying matrix elements expression constructed in frame of helicity amplitude method, if interacting energy is larger then 0.5 TeV.

The differential cross section of τ leptons production have the typical as e^- and μ^- production form but its value is significantly smaller (see, for example, fig. 2). At fig. 14 the interesting feature of total cross section dependence on energy of interacting particles is presented. The value of total cross section of two τ leptons pairs photoproduction is less then cross section of processes with two different leptons pairs photoproduction ($\gamma\gamma \rightarrow ee\mu\mu$, $\gamma\gamma \rightarrow ee\tau\tau$ and $\gamma\gamma \rightarrow \mu\mu\tau\tau$) at energy up to 150 GeV. The electrons and muons cross sections, for example, have the different behavior (see figs. 14,15).

The Table. The total cross sections dependencies on energy of initial particles. Here the following notation (α, β) is used to describe kinematical cuts, where α is the minimal angle between any two final particles, β – the minimal polar angle. Minimal admissible energy of any final particle is 1 GeV.

cut	$(3^\circ, 7^\circ)$		
energy (Gev)	$e^+e^-e^+e^-$	$\mu^+\mu^-\mu^+\mu^-$	$\tau^+\tau^-\tau^+\tau^-$
60	1254.36 ± 6.73	1210.99 ± 8.90	269.98 ± 2.82
120	379.86 ± 1.87	370.37 ± 2.48	179.01 ± 1.52
200	154.94 ± 0.81	152.87 ± 1.25	93.85 ± 0.63
300	76.41 ± 0.47	75.36 ± 0.44	52.45 ± 0.35
400	46.97 ± 0.30	46.67 ± 0.35	34.46 ± 0.30
500	32.15 ± 0.15	32.14 ± 0.27	24.15 ± 0.15
1000	9.90 ± 0.07	9.97 ± 0.12	8.35 ± 0.08
1500	5.05 ± 0.04	5.07 ± 0.11	4.31 ± 0.03
2000	3.04 ± 0.03	3.03 ± 0.03	2.66 ± 0.02
	$e^+e^-\mu^+\mu^-$	$e^+e^-\tau^+\tau^-$	$\mu^+\mu^-\tau^+\tau^-$
60	630.20 ± 5.49	354.83 ± 2.29	341.28 ± 1.42
120	189.41 ± 1.55	139.00 ± 1.09	133.51 ± 0.90
200	78.12 ± 0.54	62.80 ± 0.46	62.31 ± 0.52
300	38.77 ± 0.23	32.52 ± 0.23	32.40 ± 0.27
400	23.95 ± 0.17	20.52 ± 0.16	20.49 ± 0.12
500	16.43 ± 0.14	14.18 ± 0.12	14.15 ± 0.13
1000	5.03 ± 0.04	4.65 ± 0.04	4.56 ± 0.03
1500	2.55 ± 0.01	2.33 ± 0.02	2.30 ± 0.01
2000	1.52 ± 0.01	1.42 ± 0.01	1.40 ± 0.01




Fig.2. The differential cross section dependence of $\gamma\gamma \rightarrow e^+e^-e^+e^-$ (solid line) and $\gamma\gamma \rightarrow \tau^+\tau^-\tau^+\tau^-$ (dashed line) processes on cosine of polar angle at averaged polarization state of interacting particles. The energy of $\gamma\gamma$ - beam is 120 GeV in c.m.s.










Fig.3. The differential cross section of $\gamma\gamma \rightarrow e^+e^-\mu^+\mu^-$ process dependence on cosine of polar angle at averaged polarization state of interacting particles.



Figs.4-9. The dependence of the $\gamma\gamma \rightarrow e^+e^-e^+e^-$ differential cross section on cosine of polar angle at fixed spin states of interacting particles. For identification of particle polarization state the following notation is used: $(+, -, +, -, +, -) = (\lambda_1, \lambda_2, \lambda_3, \lambda_4, \lambda_5, \lambda_6)$, where $\lambda_{1(2)}$ corresponds to polarization state of photon with four momentum $k_{1(2)}$, $\lambda_{3,4,5,6}$ — helicity of lepton with four momentum $p_{1,2,3,4}$. Notations of $(+, -)$ and $(+, +)$ correspond to cross sections of unpolarized leptons production by scattering of circular polarized photon beams. The energy of $\gamma\gamma$ - beam is 120 GeV in c.m.s.




Fig.10. The differential cross section of $\gamma\gamma \rightarrow e^+e^-e^+e^-$ process dependence on x . Here $x = \frac{(e_1 k_1)}{s/2}$ is the scalar product of four momenta of initial photon and final positron normalized by invariant mass of incoming photons pair.




Fig.11. The differential cross section of $\gamma\gamma \rightarrow e^+e^-e^+e^-$ process dependence on x . Here $x = \frac{(e_1 e_2)}{s/2}$ is the scalar product of four momenta of final electron and final positron normalized by invariant mass of incoming photons pair.




Fig.12. The differential cross section of $\gamma\gamma \rightarrow e^+e^-e^+e^-$ process dependence on x . Here $x = \frac{E_1 + E_2}{\sqrt{s}/2}$ is the energy of any final electron-positron pair normalized by energy of incoming photons pair.




Fig.13. The differential cross section of $\gamma\gamma \rightarrow e^+e^-e^+e^-$ process dependence on x . Here $x = \frac{E_1}{\sqrt{s}/2}$ is the energy of any final electron normalized by energy of incoming photons pair.




Fig.14. The total cross sections of $\gamma\gamma \rightarrow 4l$ processes dependence on energy of interacting particles.




Fig.15. The total cross sections of $\gamma\gamma \rightarrow 4l$ processes dependence on energy of interacting particles.

References

- [1] TESLA Technical Design Report. Part.6. The Photon Collider at TESLA. (2001). DESY-01-011E;
- [2] T. Shishkina, V. Makarenko. 2002. The investigation of spin effects in photon production with fermion pair in collision. hep-ph/0212409.
- [3] V. Makarenko, K. Moenig, T. Shishkina. // Eur. Phys. J. 2003. C32. s01. p.143;
- [4] LC- PHSM-2003-016. 6. V. Makarenko, T. Shishkina. 2003. The finite mass calculations for process. hep-ph/0310104.
- [5] T.V. Shishkina, I.A. Sotsky. (2003) *Production of four leptons in $\gamma\gamma$ - interaction*, hep-ph/0312208.
- [6] P. De Causmaecker, R. Gastmans, W. Troost, Tai Tsun Wu, Phys. Lett. Vol.B105. (1981) p.215;
- [7] P. De Causmaecker, R. Gastmans, W. Troost, Tai Tsun Wu, Nucl. Phys. Vol.B206. (1982) p.53;
- [8] F.A. Berends et al., Nucl. Phys. Vol.B206. (1982) p.61;
- [9] S. Dittmaier. (1998) *Weyl-van-der-Waerden formalism for helicity amplitudes of massive particles*, hep-ph/9805445;
- [10] D. Yu. Bardin, N. M. Shumeiko, Nucl. Phys. Vol.B127. (1977) p.242;
- [11] T. V. Kukhto (Shishkina), N. M. Shumeiko, S. I. Timoshin, J. Phys. G: Nucl. Phys. Vol.13. (1987) p.725;

